



# Ultrasound evaluation of fetal gender at 12-14 weeks

Lubusky M.<sup>1,2</sup>, Prochazka M.<sup>1</sup>, Studnickova M.<sup>1</sup>, Vomackova K.<sup>3</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, University Hospital, Olomouc, Czech Republic <sup>2</sup>Department of Medical Genetics and Fetal Medicine, University Hospital, Olomouc, Czech Republic <sup>3</sup>Department of Surgery I, University Hospital, Olomouc, Czech Republic

# **OBJECTIVE**

The aim of this work was to assess the feasibility and accuracy of fetal gender assignment by transabdominal ultrasound at 12-14 weeks of gestation.

### **METHODS**

Fetal gender assessment was performed in 1222 singleton pregnancies. In all fetuses the crown-rump length (CRL) was measured and the genital area of the fetus was examined in the mid-sagittal plane. The result of ultrasound examination was compared to the phenotypic sex of the newborn after delivery.

### **RESULTS**


Feasibility as well as accuracy in determining gender increased with growing fetal CRL. Fetal gender was reliably determined (feasibility 97.4%, accuracy 100%) at CRL ≥60 mm.

At CRL ≥ 55 mm feasibility was 95.5% and accuracy 99.1% (99.8% in male gender vs 98.4% in female gender).


At CRL ≥ 50 mm the feasibility was 90.4% and accuracy 96.5%. At CRL ≥ 45 mm feasibility was 83.9% and accuracy 92.5%.

## CONCLUSION

Fetal gender may reliably be determined when CRL ≥ 60 mm (gestational age ≥ 12+2). Male gender may already be reliably determined when CRL ≥ 55 mm (gestational age ≥ 12+0). If CRL < 50 mm (gestational age < 11+4) gender cannot be reliably predicted.



**Figure 1** Ultrasound examination established gender to be **male** if the angle between the genital tubercle axis and a horizontal line through the lumbosacral skin surface was >30°.





**Figure 2** Ultrasound examination established gender to be **female** if the axis of the genital tubercle ran parallel (<10°) or convergent to the horizontal line.